



The most common electrocardiographic abnormalities in patients with diabetes mellitus

Kolsoum Teimouri¹ , Samaneh Pakravan² , Kamran Azadbakht^{3*} 

Abstract

Diabetes mellitus (DM) is a term used to describe a group of metabolic illnesses caused by an increase in blood sugar (hyperglycemia), which causes defects in insulin secretion, insulin function, or both. Cardiovascular diseases (CVDs) are the underlying cause of death in people with DM. Electrocardiography is a non-invasive clinical screening method for the early diagnosis of CVDs. This review study was performed with the aim of electrocardiogram (ECG) changes in diabetic patients.

Keywords: Cardiovascular diseases, Electrocardiography, Diabetes Mellitus

Please cite this paper as: Teimouri K, Pakravan S, Azadbakht K. The most common electrocardiographic abnormalities in patients with diabetes mellitus. *J Parathyr Dis.* 2022;10:e9142.

Copyright © 2022 The Author(s); Published by Nickan Research Institute. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Diabetes mellitus (DM) is a term used to describe a group of metabolic illnesses caused by an increase in blood sugar (hyperglycemia), which causes defects in insulin secretion, insulin function, or both (1).

The American Diabetes Association (ADA) divides DM into three general types: type 1 diabetes (insulin-dependent DM), type 2 diabetes (non-insulin-dependent DM), and gestational diabetes (2). DM is a major rapidly expanding public health problem in low- and middle-income nations (3). Diabetic patients have surged in number globally in recent decades, from 108 million in 1980 to 463 million in 2019. According to current projections, this number will approach 700 million by 2045 (4). This disease has become a major issue in most Asian nations, particularly Iran, where it is anticipated that 9.2 million people would have diabetes by 2030 (5). DM damages several organs of the body over time, including the heart, blood vessels, eyes, kidneys, and nerves; the incidence of cardiovascular is higher in diabetic patients than in the general population (6). Cardiovascular diseases (CVDs) are the leading cause of death in those who have type 2 DM (T2DM) (7). DM leads to damage to the cardiovascular system through oxidative stress, endothelial dysfunction, and atherogenesis (8). There is a link between diabetes and the risk of heart failure, which increases more than twice in males and more than five times in women, according to studies (9). Einarson et al according to a meta-analysis of medical data on 454,9481 people with T2DM showed

that nearly one-third of patients had CVDs and half of all deaths were due to CVDs (10). Electrocardiogram (ECG) is a common and non-invasive screening tool for assessing CVDs (7). Electrocardiogram is a record of electrical currents caused by stimulation of the heart muscle, there are three main components to the Electrocardiography, namely P, QRS complex, and T waves (11). In patients with DM, the ECG changes (12). The QT interval on the ECG represents the entire time of ventricular depolarization and repolarization and varies with heart rate, thus more accurately expressed as corrected QT interval (QTc), QT interval prolongation (greater than 440 milliseconds) is more common in patients with DM (13). This article aims to investigate ECG changes in patients with DM.

Materials and Methods

In this article, we review recently published works focusing on ECG changes due to DM. Articles were searched in PubMed, Google Scholar, and ScienceDirect databases with related keywords including diabetes mellitus, electrocardiography, and cardiovascular diseases.

Electrocardiographic changes

Sardesai et al, to determine ECG abnormalities examined 130 patients (58.5% male, 41.5% female) with T2DM without a history of heart disease with a mean age of 56.3 years in India. Findings of this study show that 53.1% of patients had abnormal ECG, In 17.7% of patients, changes in ST-segment and T wave occurred (14). In a

Received: 18 June 2022, Accepted: 28 July 2022, ePublished: 4 August 2022

¹Department of Medical Physics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran. ²Department of Clinical Biochemistry, Faculty of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran. ³School of Dentistry, Khorramabad University of Medical Sciences, Lorestan, Iran.

*Corresponding author: Kamran Azadbakht, Email: dr.kamran3519@gmail.com, azadbakht.k@lums.ac.ir

■ Implication for health policy/practice/research/medical education

Electrocardiographic abnormalities are common in diabetic patients. The most common ECG abnormality was QT interval prolongation.

study of 258 patients (133 males and 125 females) with T2DM in Ethiopia, Sinamaw et al found that 45% of patients underwent ECG changes. The most common ECG abnormalities included T-wave inversion (14.73%), ST-Segment depression (6.6%), prolongation of the QT interval (3.1%), and prolongation of the QRS complex duration (4.7%) (15). Charamba et al, at a clinical center in Ireland conducted a study on seventeen men with type 1 diabetes mellitus (T1DM) with a mean age of 52.5 years. They showed that mean QTc was significantly longer in these patients (16).

In a study of 100 patients with T2DM who had no history of heart disease (mean age of 50.3 years) and 50 healthy people as a control group, Gupta et al examined ECG alterations. ECG abnormalities were detected in 26% of diabetic patients, with ST-segment depression and T-wave inversion being the most prevalent, followed by left atrial enlargement (17).

In a study of 100 patients (43 males and 57 females) with T2DM with a mean age of 58.3 years in Senegal, Ndour et al Found that QT interval increased in 21% of patients, Arrhythmia was also observed in 20% of patients (18). In a case-control study in Denmark, Isaksen et al examined 855 people with T1DM (442 males, 413 females) with a mean age of 48 years without heart disease, and 1710 people as a control group were studied. They discovered that patients with T1DM had a longer QTc interval and a longer QRS complex than the control group (97 ± 11 versus 95 ± 11 ms) (12). Ninkovic et al in a study of 501 patients (227 males, 274 females) with T2DM with a mean age of 60.4 years discovered that QTc interval prolongation was significant in these patients (more than 440 ms) (13).

In a study conducted by Fagher et al in Sweden, 54% of 70 T2DM patients with an average age of 72 years died after three years of follow-up. They found that QT interval prolongation (more than 440 milliseconds) was found in 51.4% of patients and was highly linked to three-year death (19). Cox et al studied 1020 people with T2DM (569 females, 451 males) with a mean age of 61.4 years. Their study reported a prolongation of the QTc interval (more than 450 ms) (20). 35 women with gestational DM, with an average age of 32 years, and 31 healthy pregnant women were studied in the Czech Republic by Žákovičová et al (21). They found that the QT interval was significantly longer in Pregnant women with diabetes (21). In a study by Medová et al in the Czech Republic, an ECG test was taken from 3 groups of women, including group 1, 26 pregnant women with diabetes, group 2, 54 healthy pregnant women, and group 3, 18 healthy non-pregnant women. The results of this study showed that in pregnant patients with DM the duration of the QRS complex was shorter than in the other two groups (82 ± 6.8 ms(in pregnant women with diabetes, (89.5 ± 8.2 ms) in healthy pregnant women, and (90.8 ± 7.9 ms) in the control group (22). In a large study, Pfister et al analyzed 5238 people with T2DM and found that the QTc interval was increased in these patients, which was associated with heart failure (23). The results of these studies are summarized in Table 1.

Conclusion

ECG abnormalities are common in diabetic patients. The most common ECG abnormality was QT interval prolongation. Routine ECG screening should be performed for all patients with diabetes to lessen the risk of ECG abnormalities and other problems.

Authors' contribution

Conceptualization: KT and SP. Methodology: KT and SP. Investigation: KT, SP, KA and ER. Resources: KT, SP, KA and ER. Data Curation: KT, SP, KA and ER. Writing—Original Draft Preparation:

Table1. Summary of ECG findings in patients with diabetes mellitus

First author	Year	ECG findings
Sardesai et al	2022	ST-T changes
Sinamaw et al	2022	T-wave inversion, QT interval prolongation, ST-Segment Depression, prolonging QRS complex duration
Charamba et al	2021	QTc interval prolongation
Isaksen et al	2018	QTc interval prolongation, prolonging QRS complex duration
Gupta et al	2017	ST-Segment Depression, T-wave inversion
Ndour et al	2017	QT interval prolongation
Ninkovic et al	2016	QTc interval prolongation
Fagher et al	2015	QT interval prolongation
Cox et al	2014	QTc interval prolongation
Žákovičová et al	2014	QT interval prolongation
Medová et al	2012	Shortened QRS complex duration
Pfister et al	2011	QTc interval prolongation

KT, SP, KA and ER. Writing—Review and Editing: KT, SP, KA and ER. Supervision: KT, SP, KA and ER. Project Administration: KT, SP, KA and ER.

Conflicts of interest

There are no conflicts of interest declared by the authors.

Ethical issues

Ethical concerns (including plagiarism, data fabrication, and double publication) are fully respected by the authors.

Funding/Support

No financing from any source.

References

- Petersmann A, Müller-Wieland D, Müller UA, Landgraf R, Nauck M, Freckmann G, et al. Definition, classification and diagnosis of diabetes mellitus. *Exp Clin Endocrinol Diabetes*. 2019;127:S1-S7. doi: 10.1055/a-1018-9078.
- Addendum. 2. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes-2021. *Diabetes Care* 2021;44:S15-S33. doi: 10.2337/dc21-S002.
- Seiglie JA, Marcus ME, Ebert C, Prodromidis N, Geldsetzer P, Theilmann M, et al. Diabetes prevalence and its relationship with education, wealth, and BMI in 29 low- and middle-income countries. *Diabetes Care*. 2020;43:767-75. doi: 10.2337/dc19-1782.
- Cheng HT, Xu X, Lim PS, Hung KY. Worldwide Epidemiology of diabetes-related end-stage renal disease, 2000-2015. *Diabetes Care*. 2021;44:89-97. doi: 10.2337/dc20-1913.
- Fana SE, Ebrahimi R, Esmaeili S, Rambod C, Namazi N, Nasli-Esfahani E, et al. Iran diabetes research study; knowledge discovery in diagnosis: a scoping review. *J Diabetes Metab Disord*. 2021;20:1807-14. doi: 10.1007/s40200-021-00843-x.
- Valensi P, Henry P, Boccarda F, Cosson E, Prevost G, Emmerich J, et al. Risk stratification and screening for coronary artery disease in asymptomatic patients with diabetes mellitus: Position paper of the French Society of Cardiology and the French-speaking Society of Diabetology. *Diabetes Metab*. 2021;47:101185. doi: 10.1016/j.diabet.2020.08.002.
- VanHoose L, Sawers Y, Loganathan R, Vacek JL, Stehno-Bittel L, Novikova L, et al. Electrocardiographic changes with the onset of diabetes and the impact of aerobic exercise training in the Zucker Diabetic Fatty (ZDF) rat. *Cardiovasc Diabetol*. 2010;9:56. doi: 10.1186/1475-2840-9-56.
- Tokatli A, Kiliçaslan F, Alis M, Yiginer O, Uzun M. Prolonged Tp-e Interval, Tp-e/QT Ratio, and Tp-e/QTc Ratio in Patients with Type 2 Diabetes Mellitus. *Endocrinol Metab (Seoul)*. 2016;31:105-12. doi: 10.3803/EnM.2016.31.1.105.
- Rosano GM, Vitale C, Seferovic P. Heart Failure in Patients with Diabetes Mellitus. *Card Fail Rev*. 2017;3:52-5. doi: 10.15420/cfr.2016:20:2.
- Einarson TR, Acs A, Ludwig C, Panton UH. Prevalence of cardiovascular disease in type 2 diabetes: a systematic literature review of scientific evidence from across the world in 2007-2017. *Cardiovasc Diabetol*. 2018;17:83. doi: 10.1186/s12933-018-0728-6.
- Martis RJ, Acharya UR, Mandana KM, Ray AK, Chakraborty C. Application of principal component analysis to ECG signals for automated diagnosis of cardiac health. *Expert Syst Appl*. 2012;39:11792-800. doi: 10.1016/j.eswa.2012.04.072.
- Isaksen JL, Graff C, Ellervik C, Jensen JS, Rossing P, Kanter JK, et al. Cardiac repolarization and depolarization in people with Type 1 diabetes with normal ejection fraction and without known heart disease: a case-control study. *Diabet Med*. 2018;35:1337-44. doi: 10.1111/dme.13689.
- Ninkovic VM, Ninkovic SM, Miloradovic V, Stanojevic D, Babic M, Giga V, et al. Prevalence and risk factors for prolonged QT interval and QT dispersion in patients with type 2 diabetes. *Acta Diabetol*. 2016;53:737-44. doi: 10.1007/s00592-016-0864-y.
- Sardesai VV, Kokane HT, Mukherjee S, Sangle SA. A study of electrocardiographic and 2D echocardiographic changes in type 2 diabetes mellitus patients without cardiovascular symptoms. *J Family Med Prim Care*. 2022;11:1036-9. doi: 10.4103/jfmpc.jfmpc_1195_21.
- Sinamaw D, Getnet M, Abdulkadir M, Abebaw K, Ebrahim M, Diress M, et al. Patterns and associated factors of electrocardiographic abnormality among type 2 diabetic patients in Amhara National Regional State Referral Hospitals, Ethiopia: a multicenter institution-based cross-sectional study. *BMC Cardiovasc Disord*. 2022;22:230. doi: 10.1186/s12872-022-02661-2.
- Charamba B, Liew A, Coen E, Newell J, O'Brien T, Wijns W, et al. Modelling the relationship between continuously measured glucose and electrocardiographic data in adults with type 1 diabetes mellitus. *Endocrinol Diabetes Metab*. 2021;4:e00263. doi: 10.1002/edm2.263.
- Gupta S, Gupta RK, Kulshrestha M, Chaudhary RR. Evaluation of ECG abnormalities in patients with asymptomatic type 2 diabetes mellitus. *J Clin Diagn Res*. 2017;11:Oc39-41. doi: 10.7860/jcdr/2017/24882.9740.
- Ndour MA, Abdoulaye L, Codou FB, Mohamed LY, Nafy N, Diaba DN, et al. Electrocardiographic changes in patients with Type 2 diabetes multicentric cross sectional and descriptive study in Dakar. *Open J Intern Med*. 2017;7:64. doi: 10.4236/ojim.2017.73007.
- Fagher K, Nilsson A, Löndahl M. Heart rate-corrected QT interval prolongation as a prognostic marker for 3-year survival in people with Type 2 diabetes undergoing above-ankle amputation. *Diabet Med*. 2015;32:679-85. doi: 10.1111/dme.12632.
- Cox AJ, Azeem A, Yeboah J, Soliman EZ, Aggarwal SR, Bertoni AG, et al. Heart rate-corrected QT interval is an independent predictor of all-cause and cardiovascular mortality in individuals with type 2 diabetes: the Diabetes Heart Study. *Diabetes Care*. 2014;37:1454-61. doi: 10.2337/dc13-1257.
- Žákovičová E, Kittnar O, Slaviček J, Medová E, Šváb P, Charvát J. ECG body surface mapping in patients with gestational diabetes mellitus and optimal metabolic compensation. *Physiol Res*. 2014;63:S479-87. doi: 10.33549/physiolres.932921.
- Medová E, Fialová E, Mlček M, Slaviček J, Dohnalová A, Charvát J, et al. QT dispersion and electrocardiographic changes in women with gestational diabetes mellitus. *Physiol Res*. 2012;61:S49-55. doi: 10.33549/physiolres.932419.
- Pfister R, Cairns R, Erdmann E, Schneider CA. Prognostic impact of electrocardiographic signs in patients with Type 2 diabetes and cardiovascular disease: results from the PROactive study. *Diabet Med*. 2011;28:1206-12. doi: 10.1111/j.1464-5491.2011.03281.x.